


- born 1917 in Freiburg, Germany
- studied Chemistry in Munich till 1938 (diploma), 1940 (PhD)
- 1938-1946 private assistant of Prof. Wieland
- 1948-1950 lecturer at the Harvard University
- US citizenship in 1953
- from 1957 Chief of the Laboratory of Chemistry, National Institute of Arthritis and Metabolic Diseases, Bethesda, MD
- his research interests were versatile, isolation of natural products, reaction mechanisms, synthetic organic chemistry, and biochemistry
- more than 300 publications, many of them in J. Am. Chem. Soc.




Biochemistry - NIH-Shift¹

expected reaction



observed reactions




mechanism?

¹ G. Gruoff, J. W. Daly, D. M. Jerina, J. Renson, B. Witkop, S. Udenfriend, *Science* **1967**, 157, 1524-1530.


Biochemistry- protein structure²

- Enzymatic and non-enzymatic pathways to cleave amide bonds selectively were important to determine the primary sequence and the reactive center of enzymes
- cyanogen bromide cleaves the amide bond next to methionine




- Only a slow reaction with cysteine, no reaction with all the other amino acids

- NBS in 8.0 M urea cleaves amide bonds next to tryptophanes




- No side reactions with tyrosine or other aromatic amino acids
- interesting reaction at pH above pH 9




² B. Witkop, *Science* **1968**, 162, 318-326.

Isolation and determination of Natural Products

structure of yohimbine³


- The general structure was known, but not the absolute stereo-chemistry of the stereocenters at C15 and C20
- treatment of erectile dysfunction
- Increases blood flow in extremities
- many side effects!! (high blood pressure, anxiety.....)




- by comparison of IR-data and optical activity of synthetic and the isolated decahydro-isoquinoline they could figure out the absolute stereo-chemistry of C15 and C20

³ B. Witkop, *J. Am. Chem. Soc.* **1949**, 71, 2559-2566.

Photochemistry


Chemistry with thymine dimers⁴

easy access to highly substituted
diastereomerically pure cyclobutanes


Photochemistry also with
3-deazapyrimidone⁵

⁴ T. Kunieda, B. Witkop, *J. Am. Chem. Soc.* **1971**, 93, 3493-3499.


⁵ B. Witkop *et al.*, *Lieb. Ann. Chem.* **1988**, 979-981.

Photochemistry – cage compounds⁶⁶ B. Witkop *et al.*, *J. Am. Chem. Soc.* **1974**, 96, 2564-2569.

Photochemistry

Stereochemistry of photocyclisations with heterocyclic anilides⁷

Explain the different mechanisms leading to the *cis* and the *trans* product!?


Photooxidation of Trp to kynurenine⁸

⁷ B. Witkop *et al.*, *J. Org. Chem.* **1975**, 40, 3001-3003.


⁸ B. Witkop *et al.*, *Proc. Natl. Acad. Sci.* **1977**, 4730-4732.

Synthesis and Mechanisms

Living on the edge....⁹



Fischer indole synthesis in polyphosphoric acid¹⁰




Good yields, easy workup

Limitation: only the synthesis of 2-substituted indoles possible

⁹ B. Witkop *et al.*, *J. Am. Chem. Soc.* **1969**, 91, 6199-6200.

¹⁰ B. Witkop *et al.*, *J. Am. Chem. Soc.* **1952**, 74, 3948-3949.


Synthesis and Mechanisms - Chemistry of Spirooxindoles






¹¹ J. B. Patrick, B. Witkop, *J. Am. Chem. Soc.* **1952**, 75, 2572-2576.

Synthesis and Mechanisms - Twofold Wagner-Meerwein-Rearrangements¹²⁻¹³

Expected reaction with Lithiumorganyl



Unexpected reaction with Grignard reagent





Rearrangement can be induced by an excess Grignard reagent, acid or BF_3OEt_2

¹² J. B. Patrick, B. Witkop, *J. Am. Chem. Soc.* **1951**, 73, 1558-1564.

¹³ A. Ek, B. Witkop, *J. Am. Chem. Soc.* **1951**, 73, 5664-5669.


Synthesis and Mechanisms

Deketopiperazines of 3,4-dehydroproline – unique structural properties¹⁴Stereodiagram of the *L,L*-diketopiperazine¹⁴ B. Witkop *et al.*, *J. Am. Chem. Soc.* **1972**, 96, 539-543.

Synthesis and Mechanisms


Batrachotoxin¹⁵

- a cardiotoxic alkaloid from the colombian arrow poison frog *Phylllobates aurotaenia* ($LD_{50} = 2\mu\text{g}/\text{kg}$ mice)
- very labil venom (4 expeditions to Colombia were necessary) and most of the experiments were carried out in μg -quantities



- by NMR, UV/Vis and MS they found that R is 2,4-dimethylpyrrole-3-carboxylic acid

- ultimate proof of their hypothesis was the partial synthesis of Batrachotoxine from Batrachotoxine A and activated 2,4-dimethylpyrrole-3-carboxylic acid



¹² T. Tokuyama, J. Daly, B. Witkop, *J. Am. Chem. Soc.* **1969**, 91, 3931-3938.