

1. Electronic absorption band assignments for copper (II) salicylaldimine complexes. T.N. Waters and P.E. Wright (1971) *J. Inorg. Nucl. Chem.* **33**, 359-363.
2. Conformational influences in copper co-ordination compounds. Part V. Crystal and molecular structure of [1,2-bis(2-aminobenzylidene)-amino]propanato(2-) copper (II). D. Hall, T.N. Waters and P.E. Wright (1973) *J. Chem. Soc. Dalton*, 1508-1512.
3. Outline structure of cytochrome c_3 and consideration of its properties. C.M. Dobson, N.J. Hoyle, C.F. Gerald, M. Bruschi, J. LeGall, P.E. Wright and R.J.P. Williams (1974) *Nature* **249**, 425-429.
4. E.S.R. and optical spectral properties of copper (II) complexes with Schiff base ligands derived from o-aminobenzaldehyde. G.A. Bowmaker, T.N. Waters and P.E. Wright (1975) *J. Chem. Soc. Dalton*, 867-870.
5. Electronic transitions in salicylaldimine complexes. A.C. Braithwaite, P.E. Wright and T.N. Waters (1975) *J. Inorg. Nucl. Chem.* **37**, 1669-1674.
6. Assignment of NMR spectrum of iron (III) protoporphyrin (IX) dicyanide using paramagnetic shift and broadening probes. J.G. Brassington, R.J.P. Williams and P.E. Wright (1975) *J.C.S. Chem. Commun.*, 338-340.
7. Conformational studies of peroxidase-substrate complexes. Structure of the indolepropionic acid-horseradish peroxidase complex. P.S. Burns, R.J.P. Williams and P.E. Wright (1975) *J.C.S. Chem. Commun.*, 795-796.
8. Proton magnetic resonance studies of peroxidases from turnip and horseradish. R.J.P. Williams, P.E. Wright, G. Mazza and J.R. Ricard (1975) *Biochim. Biophys. Acta* **412**, 127-147.
9. Pulse methods for the simplification of protein NMR spectra. I.D. Campbell, C.M. Dobson, R.J.P. Williams and P.E. Wright (1975) *FEBS Lett.* **57**, 96-99.
10. A study of redox reactions of biological importance between Fe (III) complexes and aromatic moieties. P.S. Burns, J.F. Harrod, R.J.P. Williams and P.E. Wright (1976) *Biochim. Biophys. Acta* **428**, 261-268.
11. Nuclear magnetic resonance studies of the adrenal gland and some other organs. A. Daniels, R.J.P. Williams and P.E. Wright (1976) *Nature* **261**, 321-322.
12. High resolution NMR studies of soybean leghemoglobin *a*. P.E. Wright and C.A. Appleby (1977) *FEBS Lett.* **78**, 61-66.
13. Oxidation and reduction properties of cytochromes and peroxidases. R.J.P. Williams, G.R. Moore and P.E. Wright, (1977) *In: Biological Aspects of Inorganic Chemistry.*, Eds. A.W. Addison, W.R. Cullen, D. Dolphin and B.R. James, Wiley, New York, 369-401.
14. The proton NMR spectra of whole organs. A. Daniels, J. Krebs, B.A. Levine, P.E. Wright and R.J.P. Williams, (1977) *In: NMR in Biology*, Eds. R.A. Dwek, I.D. Campbell, R.E. Richards and R.J.P. Williams, Academic Press, 277-287.
15. The storage of small molecules in the octopus salivary gland: nuclear magnetic resonance study. A. Daniels, J. Krebs, P.E. Wright and R.J.P. Williams (1977) *Biochem. Soc. Trans.* **5**, 1149-1151.
16. High resolution proton magnetic resonance studies of plastocyanin. H.C. Freeman, V.A. Norris, J.A.M. Ramshaw, and P.E. Wright (1977) *In: Proceedings of the Fourth International Congress of Photosynthesis*, 805-809.
17. High resolution proton magnetic resonance studies of plastocyanin. H.C. Freeman, V.A. Norris, J.A.M. Ramshaw and P.E. Wright (1978) *FEBS Lett.* **86**, 131-135.

18. The character of stored molecules in chromaffin granules of the adrenal medulla: a nuclear magnetic resonance study. A.J. Daniels, R.J.P. Williams and P.E. Wright (1978) *Neurosci.* **3**, 573-585.
19. Spin state equilibria in soybean ferric leghemoglobin and its complexes with formate and acetate. J. Trehwella, P.E. Wright and C.A. Appleby (1979) *Biochem. Biophys. Res. Commun.* **88**, 713-721.
20. Molecular basis for proton-dependent anion binding by soybean leghemoglobin **a**. J. Trehwella, P.E. Wright and C.A. Appleby (1979) *Nature* **280**, 87-88.
21. Conformation and dynamics of biological macromolecules in solution. P.E. Wright (1979) *Proc. Aust. Soc. Biophysics* **3**, 1S-5S.
22. Electron transfer reagent binding sites on plastocyanin. D.J. Cookson, M.T. Hayes and P.E. Wright (1980) *Nature* **283**, 682-683.
23. NMR study of the interactions of plastocyanin with chromium (III) analogues of inorganic electron transfer reagents. D.J. Cookson, M.T. Hayes and P.E. Wright (1980) *Biochim. Biophys. Acta.* **591**, 162-176.
24. ¹H NMR and EPR studies on skeletal muscle actin indicate that the metal and nucleotide binding sites are separate. J.A. Barden, R. Cooke, P.E. Wright and C.G. dos Remedios (1980) *Biochemistry* **19**, 5912-5916.
25. ¹H NMR studies of ferric soybean leghemoglobin. Assignment of hyperfine shifted resonances of complexes with cyanide, nictinate, pyridine and azide. J. Trehwella and P.E. Wright (1980) *Biochim. Biophys. Acta* **625**, 202-220.
26. Resonance Raman evidence for constrained heme structure in soybean leghemoglobin and its derivatives. R.A. Armstrong, M.J. Irwin and P.E. Wright (1980) *Biochem. Biophys. Res. Commun.* **95**, 682-689.
27. Measurement of heme accessibility in soybean ferric leghemoglobin *a* and its complexes by proton magnetic relaxation. D.L. Ollis, P.E. Wright, J.M. Pope and C.A. Appleby (1981) *Biochemistry* **20**, 587-594.
28. Heme pocket structure, conformational mobility and heme accessibility of soybean leghemoglobin. P.E. Wright and C.A. Appleby, 1981 in *Current Perspectives in Nitrogen Fixation*, Eds. A.H. Gibson and W.E. Newton, Australian Academy of Science, 365.
29. Resonance Raman studies of soybean leghemoglobin and myoglobin. Origin of the differences in O₂ dissociation rate constants. M.J. Irwin, R.S. Armstrong and P.E. Wright (1981) *FEBS Lett.* **133**, 239-243.
30. Molecular basis for the pH-dependent binding of O₂ to soybean leghemoglobin. P.E. Wright and C.A. Appleby (1981) *Proc. Aust. Biochem.* **14**, 33.
31. Soret excited resonance Raman spectrum of carbonmonoxyleghemoglobin: assignment of $\square_{\text{Fe-CO}}$. R.S. Armstrong, M.J. Irwin and P.E. Wright (1982) *J. Amer. Chem. Soc.* **104**, 626-627.
32. Differences in the heme environment of soybean leghemoglobin components shown by ¹H NMR spectroscopy. C.A. Appleby, J. Trehwella and P.E. Wright (1982) *Biochim. Biophys. Acta* **700**, 171-177.
33. The mobile distal histidine of leghemoglobin: Does it control oxygen binding kinetics? C.A. Appleby, W.E. Blumberg, J.H. Bradbury, W.H. Fuchsman, J. Peisach, B.A. Wittenberg, J.B. Wittenberg and P.E. Wright (1982) *In: Interactions Between Iron and Proteins in Oxygen and Electron Transport*, Ed. C. Ho, Elsevier, 435-441.

34. A comment on resonance Raman spectroscopic investigations of axial ligation of the heme iron atom in soybean leghemoglobin *a*. R.S. Armstrong, M.J. Irwin, J.E. Wellington and P.E. Wright (1982) *Acta Chem. Scand.* **36B**, 263-265.
35. A two-dimensional NMR method for assignment of imidazole ring proton resonances of histidine residues in proteins. G. King and P.E. Wright (1982) *Biochem. Biophys. Res. Commun.* **106**, 559-565.
36. ¹³C NMR studies of chlorophyll biosynthesis in higher plants: an unequivocal proof of the participation of the C₅ pathway and evidence of a new route for the incorporation of glycine. R.J. Porra, O. Klein and P.E. Wright (1982) *Biochem. Internat.* **5**, 345-350.
37. ¹H NMR studies of complexes of ferric leghemoglobin with substituted pyridines and nicotinic acids. B.C. Mabbutt and P.E. Wright (1983) *J. Inorg. Biochem.* **18**, 123-132.
38. Leghemoglobin. Kinetic, NMR and optical studies of pH dependence of oxygen and carbon monoxide binding. C.A. Appleby, J.H. Bradbury, R.J. Morris, B.A. Wittenberg, J.B. Wittenberg and P.E. Wright (1983) *J. Biol. Chem.* **258**, 2254-2259.
39. Assignment of heme and distal amino acid proton resonances in the NMR spectra of the oxygen and carbonmonoxide complexes of soybean leghemoglobin. B.C. Mabbutt and P.E. Wright (1983) *Biochim. Biophys. Acta* **744**, 281-290.
40. A proton NMR study of the conformation of Trp-Gly-Ala-Glu in dimethylsulphoxide. G.R. Beilharz, P. Mack, A.V. Robertson and P.E. Wright (1983) *Aust. J. Chem.* **36**, 751-758.
41. NMR studies of oxyleghemoglobin. Assignment of distal histidine proton resonances and evidence for pH-dependent changes in conformation. B.C. Mabbutt, C.A. Appleby and P.E. Wright (1983) *Biochim. Biophys. Acta* **749**, 281-288.
42. Applications of two-dimensional relayed coherence transfer experiments to ¹H NMR studies of macromolecules. G. King and P.E. Wright (1983) *J. Mag. Res.* **54**, 328-332.
43. The proof by ¹³C NMR spectroscopy of the predominance of the C₅ pathway over the Shemin pathway in chlorophyll biosynthesis in higher plants and of the formation of the methyl ester group of chlorophyll from glycine. R.J. Porra, O. Klein and P.E. Wright (1983) *Eur. J. Biochem.* **130**, 509-516.
44. An examination of the basic blue copper protein from cucumber peelings by ¹H NMR. G. King, T.A. Andary, H.C. Freeman, L. Gavrilovic and P.E. Wright (1984) *FEBS Lett.* **166**, 288-292.
45. Unequivocal proof of chlorophyll formation by the C₅ pathway and tentative evidence for Shemin pathway activity in greening leaves. Is mitochondrial haem formed by the Shemin pathway in leaf cells? R.J. Porra, O. Klein and P.E. Wright (1984), In: *Advances in Photosynthesis Research*, Eds. C. Sybesma, M. Nijhoff, and W. Junk, Vol. IV, 6.725-6.728.
46. NMR studies of myelin basic protein - XI. Conformation of a nona-peptide from a segment encephalitogenic in Rhesus monkey. G.L. Mendz, W.J. Moore, G. King and P.E. Wright (1984) *Int. J. Peptide Protein Chem.* **24**, 208-217.
47. Assignment of acyl chain resonances from membranes of mammalian cells by two dimensional methods. K.J. Cross, K.T. Holmes, C.E. Mountford and P.E. Wright (1984) *Biochemistry* **23**, 5895-5897.
48. NMR and kinetic characterization of the interaction between French bean plastocyanin and horse cytochrome *c*. G.C. King, R.A. Binstead and P.E. Wright (1985) *Biochim. Biophys. Acta* **806**, 262-271.

49. ^1H NMR studies of the conformation of eosinophil chemotactic tetra-peptides and analogues in DMSO. G.R. Beilharz, J.A. Smith, K.F. Austen and P.E. Wright (1985) *Int. J. Peptide Protein Chem.* **25**, 337-346.
50. NMR studies of the structure and dynamics of monomeric hemoglobins and myoglobins. P.E. Wright, R.M. Cooke, K.J. Cross, B.C. Mabbutt, B.A. Messerle and J.E. Wellington (1985) *In: Magnetic Resonance in Biology and Medicine*, Eds. G. Govil, C.L. Khetrpal and A. Saran, Tata-McGraw-Hill, New Delhi, 131-150.
51. Calibration of ring current models for the heme ring. K.J. Cross and P.E. Wright (1985) *J. Magn. Res.* **64**, 220-231.
52. Acid dissociation constants for plastocyanin in the Cu(I) state. J.D. Sinclair-Day, M.J. Sisley, A.G. Sykes, G.C. King and P.E. Wright (1985) *Chem. Comm.* 505-507.
53. Conformational disorder of the distal leucine in monomeric *Glycera* hemoglobins and implications for oxygen binding. R.M. Cooke and P.E. Wright (1985) *FEBS. Lett.* **187**, 219-223.
54. Completely synthetic vaccine effective against heterologous enterotoxigenic *Escherichia coli*. R.A. van Houghten, R.A. Lerner, S.R. Hoffman, P.A. Worrell, P.E. Wright and F.A. Klipstein (1985), *In: Vaccines '85. Molecular and Chemical Basis of Resistance to Parasitic, Bacterial and Viral Diseases*, Eds. R.A. Lerner, R.M. Chanock and F. Brown, Cold Spring Harbor Laboratory, New York, 91-94.
55. Simplification of ^1H NMR spectra of proteins by one-dimensional multiple quantum filtration. M. Rance, C. Dalvit and P.E. Wright (1985) *Biochem. Biophys. Res. Commun.* **131**, 1094-1102.
56. Assignment of heme and distal amino acid resonances in the ^1H NMR spectra of the carbon monoxide and oxygen complexes of sperm whale myoglobin. B.C. Mabbutt and P.E. Wright (1985) *Biochim. Biophys. Acta* **832**, 175-185.
57. The immunodominant site of a synthetic immunogen has a conformational preference in water for a type II reverse turn. H.J. Dyson, K.J. Cross, R.A. Houghten, I.A. Wilson, P.E. Wright and R.A. Lerner (1985) *Nature* **318**, 480-483.
58. Antipeptide antibodies and the disorder-order phenomenon. P.E. Wright, H.J. Dyson, M. Rance, J. Ostresh, R.A. Houghten, I.A. Wilson and R.A. Lerner (1986), *In: New Approaches to Immunization*, Eds. R.M. Chanock, F. Brown and R.A. Lerner, Cold Spring Harbor Press, New York, 15-19.
59. Differences in amino acid composition and heme electronic structure of the multiple monomeric hemoglobin components of *Glycera dibranchiata*. R.M. Cooke and P.E. Wright (1985) *Biochim. Biophys. Acta* **832**, 357-364.
60. Heme orientation in the major monomeric hemoglobins of *Glycera dibranchiata*. R.M. Cooke and P.E. Wright (1985) *Biochim. Biophys. Acta* **832**, 357-372.
61. Selection by site-directed antibodies of small regions of peptides which are ordered in water. H.J. Dyson, K.J. Cross, J. Ostresh, R.A. Houghten, I.A. Wilson, P.E. Wright and R.A. Lerner (1986), *In: Synthetic Peptides as Antigens*, Ciba Foundation Symposium 119, John Wiley & Sons, Chichester, pp. 58-75.
62. Analysis of ^1H NMR spectra of proteins using multiple quantum coherence. M. Rance and P.E. Wright (1986) *J. Magn. Res.* **66**, 372-378.
63. ^1H NMR studies of high spin complexes of soybean leghemoglobin. Interactions between the distal histidine and acetate, formate and fluoride ligands. J. Trewhella, C.A. Appleby and P.E. Wright (1986) *Aust. J. Chem.* **39**, 317-324.

64. Assignment of methylene resonances in NMR spectra of cancer cells to plasma membrane triglyceride. G.L. May, L.C. Wright, K.T. Holmes, P.G. Williams, I.C.P. Smith, P.E. Wright, R.M. Fox and C.E. Mountford (1986) *J. Biol. Chem.* **261**, 3048-3053.
65. Selection rule violations in multiple quantum NMR spectroscopy. M. Rance and P.E. Wright (1986) *Chem. Phys. Lett.* **124**, 572-575.
66. Proton NMR studies of plastocyanin: Assignment of aromatic and methyl group resonances from 2D spectra. G. King and P.E. Wright (1986) *Biochemistry* **25**, 2364-2374.
67. Differentiation of direct and remote connectivities in phase sensitive double quantum spectra of proteins by variation of the multiple quantum excitation period. C. Dalvit, M. Rance and P.E. Wright (1986) *J. Magn. Reson.* **69**, 356-361.
68. ¹H NMR studies of heme pocket conformation in zinc-substituted leghemoglobin, a diamagnetic analogue of deoxyhemoglobin. C. Dalvit, L. Tennant and P.E. Wright (1986) *J. Inorg. Biochem.* **28**, 303-309.
69. The order-disorder paradox in antigen-antibody union: anti-peptide antibodies as a probe for structured regions of small peptides. H.J. Dyson, M. Rance, R.A. Houghten, P.E. Wright and R.A. Lerner (1987), *In: Biological Organization: Macromolecular Interactions at High Resolution*, Eds. R.M. Burnett and H.J. Vogel, Academic Press, pp. 227-234.
70. Kinetic and mechanisms of the oxidation of myoglobin by Fe(III) and Cu(II) complexes. K. Hegetschweiler, P. Saltman, C. Dalvit and P.E. Wright (1987) *Biochim. Biophys. Acta* **912**, 384-397.
71. Detection of long-range couplings in proteins via double quantum spectroscopy: Connection of aliphatic and imidazole resonances of histidine residues. C. Dalvit, P.E. Wright and M. Rance (1987) *J. Magn. Reson.* **71**, 539-543.
72. Assignment of resonances in the ¹H NMR spectrum of the carbon monoxide complex of sperm whale myoglobin by phase sensitive two-dimensional techniques. C. Dalvit and P.E. Wright (1987) *J. Mol. Biol.* **194**, 313-327.
73. Assignment of resonances in the ¹H NMR spectrum of the carbon monoxide complex of human hemoglobin α -chains. C. Dalvit and P.E. Wright (1987) *J. Mol. Biol.* **194**, 329-339.
74. Site-selective observation of nuclear Overhauser effects in proteins via isotopic labelling. M. Rance, P.E. Wright, B.A. Messerle and L.D. Field (1987) *J. Am. Chem. Soc.* **109**, 1591-1593.
75. ¹H resonances of proximal histidine in CO complexes of hemoglobins provide a sensitive probe of coordination geometry. C. Dalvit, L. Tennant and P.E. Wright (1987) *FEBS Lett.* **213**, 289-292.
76. A modified approach to identification of ¹H spin systems in proteins. W.J. Chazin and P.E. Wright (1987) *Biopolymers* **26**, 973-977.
77. Identification of folded structures in immunogenic peptides by 2D NMR spectroscopy. P.E. Wright, H.J. Dyson, M. Rance, R.A. Houghten and R.A. Lerner (1987), *In: Protides of the Biological Fluids*, Ed. H. Peeters, Vol. 35, Pergamon Press, Oxford, pp. 477-480.
78. NMR studies of the heme pocket conformations of monomeric hemoglobins from *Glycera dibranchiata*. Implications for ligand binding. R.M. Cooke, C. Dalvit, S.S. Narula and P.E. Wright (1987) *Eur. J. Biochem.* **166**, 399-408.

79. Structural consequences of heme isomerism in monomeric hemoglobins from *Glycera dibranchiata*. R.M. Cooke and P.E. Wright (1987) *Eur. J. Biochem.* **166**, 409-414.
80. Kinetic studies on 1:1 electron-transfer reactions involving blue copper proteins. Part 15. The reactivity of *Anabaena variabilis* plastocyanin with inorganic complexes and related NMR studies. M.P. Jackman, J.D. Sinclair-Day, M.J. Sisley, A.G. Sykes, L.A. Denys and P.E. Wright (1987) *J. Amer. Chem. Soc.* **109**, 6443-6449.
81. Complete assignment of lysine resonances in ^1H NMR spectra of proteins as probes of surface structure and dynamics. W.J. Chazin, M. Rance and P.E. Wright (1987) *FEBS Lett.* **222**, 109-114.
82. Electron transfer from cytochrome b_5 to iron and copper complexes. L.S. Reid, H.B. Gray, C. Dalvit, P.E. Wright and P. Saltman (1987) *Biochemistry* **26**, 7102-7107.
83. Folding of immunogenic peptide fragments of proteins in water solution. I. Sequence requirements for the formation of a reverse turn. H.J. Dyson, M. Rance, R.A. Houghten, R.A. Lerner and P.E. Wright (1988) *J. Mol. Biol.* **201**, 161-200.
84. Folding of immunogenic peptide fragments of proteins in water solution. II. The nascent helix. H.J. Dyson, M. Rance, R.A. Houghten, P.E. Wright and R.A. Lerner (1988) *J. Mol. Biol.* **201**, 201-217.
85. The physical basis induction of protein-reactive anti-peptide antibodies. H.J. Dyson, R.A. Lerner and P.E. Wright (1988) *Ann. Rev. Biophysics Biophys. Chem.* **17**, 305-324.
86. Kinetic studies on 1:1 electron transfer reactions involving blue copper proteins.16. The reactivity of plastocyanin from the green alga *Scenedesmus obliquus* with inorganic redox partners and related NMR studies. J. McGinnis, J.D. Sinclair-Day, A.G. Sykes, R. Powls, J. Moore and P.E. Wright (1988) *Inorg. Chem.* **27**, 2306-2312.
87. Complete assignment of the ^1H NMR spectrum of French bean plastocyanin: application of an integrated approach to spin system identification in proteins. W.J. Chazin, M. Rance and P.E. Wright (1988) *J. Mol. Biol.* **202**, 603-622.
88. Complete assignment of the ^1H NMR spectrum of French bean plastocyanin: sequential resonance assignments, secondary structure and global fold. W.J. Chazin and P.E. Wright (1988) *J. Mol. Biol.* **202**, 623-636.
89. Structural differences between oxidized and reduced thioredoxin monitored by two-dimensional ^1H NMR spectroscopy. H.J. Dyson, A. Holmgren and P.E. Wright (1988) *FEBS Lett.* **228**, 254-258.
90. Three-dimensional solution structure of plastocyanin from the green alga *Scenedesmus obliquus*. J.M. Moore, D.A. Case, W.J. Chazin, G.P. Gippert, T.F. Havel, R. Powls and P.E. Wright (1988) *Science* **240**, 314-317.
91. Conformation of peptide fragments of proteins in aqueous solution: implications for initiation of protein folding. P.E. Wright, H.J. Dyson and R.A. Lerner (1988) *Biochemistry* **27**, 7167-7175.
92. ^1H NMR studies of plastocyanin from *Scenedesmus obliquus*: Complete sequence-specific assignment, secondary structure analysis and global fold. J.M. Moore, W.J. Chazin, R. Powls and P.E. Wright (1988) *Biochemistry* **27**, 7806-7816.
93. ^1H NMR studies of human C3a anaphylatoxin in solution: sequential resonance assignments, secondary structure and global fold. W.J. Chazin, T.E. Hugli and P.E. Wright (1988) *Biochemistry* **27**, 9139-9148.
94. Isotope-edited NMR studies of Fab-peptide complexes. P. Tsang, T.M. Fieser, R.A. Houghten, R.A. Lerner and P.E. Wright (1988) *Pept. Res.* **1**, 87-92.

95. NMR studies of the conformations of leghemoglobins from soybean and lupin. S.S. Narula, C. Dalvit, C.A. Appleby and P.E. Wright (1988) *Eur. J. Biochem.* **178**, 419-435.
96. Multiple Quantum NMR. M. Rance, W.J. Chazin, C. Dalvit and P.E. Wright (1989), *In: Methods in Enzymology*, Eds. N. Oppenheimer and T.L. James, Vol. 176, 114-134.
97. Chemical modification of bovine pancreatic trypsin inhibitor for single site coupling of immunogenic peptides for NMR conformational analysis. S. Ebina, R.A. Lerner and P.E. Wright (1989) *J. Biol. Chem.* **264**, 7882-7888.
98. ¹H NMR studies of the solution conformations of an analogue of the C-peptide of ribonuclease A. J.J. Osterhout, Jr., R.L. Baldwin, E.J. York, J.M. Stewart, H.J. Dyson and P.E. Wright (1989) *Biochemistry* **28**, 7059-7064.
99. Folding of peptide fragments of proteins in water solution. P.E. Wright, H.J. Dyson and R.A. Lerner (1989), *In: Protein Folding. Deciphering the Second Half of the Genetic Code*, Eds. J.A. King and L.M. Gierasch, AAAS, 95-102.
100. Assignment of the proton NMR spectrum of reduced and oxidized thioredoxin: sequence-specific assignments, secondary structure and global fold. H.J. Dyson, A. Holmgren and P.E. Wright (1989) *Biochemistry* **28**, 7074-7087.
101. Folding of peptide fragments of proteins in aqueous solution. P.E. Wright, H.J. Dyson, V.A. Feher, L. Tennant, J.P. Waltho, R.A. Lerner and D.A. Case (1990), *In: Frontiers of NMR in Molecular Biology*, UCLA Symposia on Molecular and Cellular Biology, New Series, Eds. D. Live, I. Armitage and D. Patel, Vol. 109, Alan R. Liss, Inc., New York, 1-13.
102. Solution NMR studies of Fab'-peptide complexes. P. Tsang, T.M. Fieser, J.M. Ostresh, R.A. Houghten, R.A. Lerner and P.E. Wright (1990) *In: Frontiers of NMR in Molecular Biology*, UCLA Symposia on Molecular and Cellular Biology, New Series, Eds. D. Live, I. Armitage and D. Patel, Vol. 109, Alan R. Liss, Inc., New York, 63-73.
103. Conformation of a T cell stimulating peptide in aqueous solution. J.P. Waltho, R.A. Lerner and P.E. Wright (1989) *FEBS Lett.* **250**, 400-404.
104. What can two-dimensional NMR tell us about proteins? P.E. Wright (1989) *Trends Biochem. Sci.* **14**, 255-260.
105. Three-dimensional solution structure of a single zinc finger DNA-binding domain. M.S. Lee, G.P. Gippert, K.V. Soman, D.A. Case and P.E. Wright (1989) *Science* **245**, 635-637.
106. Complete assignment of the ¹H nuclear magnetic resonance spectrum of a synthetic zinc finger from *Xfin*. Sequential resonance assignments and secondary structure. M.S. Lee, J. Cavanagh and P.E. Wright (1989) *FEBS Lett.* **254**, 159-164.
107. Folding of peptide fragments of proteins in water solution: implications for initiation of protein folding. P.E. Wright, R.A. Lerner and H.J. Dyson (1989), in *Advances in Protein Design*, Eds. H. Blocker, J. Collins, R.D. Schmid, D. Schomburg, VCH Publishers, 13-19.
108. Three-dimensional solution structure of the reduced form of *Escherichia coli* thioredoxin determined by NMR spectroscopy. H.J. Dyson, G.P. Gippert, D.A. Case, A. Holmgren and P.E. Wright (1990) *Biochemistry* **29**, 4129-4136.

109. Proton Nuclear Magnetic Resonance assignments and solution structure of a synthetic zinc finger from *Xfin*. M.S. Lee, G.P. Gippert, K.V. Soman, D.A. Case and P.E. Wright (1990), In: Proceedings of Sixth Conversation in Biomolecular Stereodynamics. *Structure and Methods: Volume 2, DNA-Protein Complexes and Proteins*, Eds. R.H. Sarma and M.H. Sarma, Adenine Press, Albany, NY, 83-91.
110. Computational methods for determining protein structures from NMR data. G.P. Gippert, P.F. Yip, P.E. Wright and D.A. Case (1990) *Biochem. Pharmacol.* **40**, 15-22.
111. The H-helix of myoglobin as a potential independent protein folding domain. J.P. Waltho, V.A. Feher and P.E. Wright (1990), In: *Current Research in Protein Chemistry: Techniques, Structure and Function*, The Protein Society, Ed. J.J. Villafranca, Academic Press, San Diego, 283-293.
112. Antigen-antibody interactions: an NMR approach. P.E. Wright, H.J. Dyson, R.A. Lerner, L. Riechmann and P. Tsang. (1990) *Biochem. Pharmacol.* **40**, 83-88.
113. Conformational preferences of synthetic peptides derived from the immunogenic site of the circumsporozoite protein of *Plasmodium falciparum* by ¹H NMR. H.J. Dyson, A.C. Satterthwait, R.A. Lerner and P.E. Wright (1990) *Biochemistry* **29**, 7828-7837.
114. Signal suppression in the frequency domain to remove undesirable resonances with dispersive lineshapes. P. Tsang, P.E. Wright and M. Rance (1990) *J. Magn. Reson.* **88**, 210-215.
115. Structural characterization of a partly folded apomyoglobin intermediate. F.M. Hughson, P.E. Wright and R.L. Baldwin (1990) *Science* **249**, 1544-1548.
116. ¹H NMR assignments of the *Escherichia coli* dihydrofolate reductase complex with folate: Evidence for a unique conformation of bound folate. C.J. Falzone, S.J. Benkovic and P.E. Wright (1990) *Biochemistry* **29**, 9667-9677.
117. Specific deuteration strategy for enhancing direct nuclear Overhauser effects in high molecular weight complexes. P. Tsang, P.E. Wright and M. Rance (1990) *J. Amer. Chem. Soc.* **112**, 8183-8185.
118. The conformational restriction of synthetic vaccines for malaria. A.C. Satterthwait, L.-C. Chiang, T. Arrhenius, E. Cabezas, F. Zavala, H.J. Dyson, P.E. Wright and R.A. Lerner (1990) *Bull. World Hlth Organiz.* **68** (Suppl), 17-25.
119. Zinc is required for folding and binding of a single zinc finger to DNA. M.S. Lee, J.M. Gottesfeld and P.E. Wright (1991) *FEBS Lett.* **279**, 289-294.
120. Sensitivity improvement in proton-detected two-dimensional heteronuclear relay spectroscopy. J. Cavanagh, A.G. Palmer III, P.E. Wright & M. Rance (1991) *J. Magn. Reson.* **91**, 429-436.
121. Isotope-edited nuclear magnetic resonance studies of Fab-peptide complexes. P. Tsang, M. Rance and P.E. Wright (1991) In: *Methods in Enzymology*, Vol. 203, 241-261.
122. The three-dimensional solution structure of human anaphylatoxin C3a. M.W. Kalnik, W.J. Chazin and P.E. Wright (1991) In: *Techniques in Protein Chemistry II*, Ed. J.J. Villafranca, Academic Press, San Diego, pp. 393-400.
123. Defining solution conformations of small linear peptides. H.J. Dyson and P.E. Wright (1991) *Ann. Rev. Biophysics Biophys. Chem.* **20**, 519-538.
124. Kinetics and mechanism of reduction of Cu(II) and Fe (III) complexes by soybean leghemoglobin *a*. D.A. Bakan, P. Saltman, Y. Theriault and P.E. Wright (1991) *Biochim. Biophys. Acta*, **1079**, 182-196

125. Mapping the anatomy of the immunodominant domain of the HIV GP41 transmembrane protein: Peptide conformation analysis using monoclonal antibodies and proton nuclear magnetic resonance spectroscopy. M.B.A. Oldstone, A. Tishon, H. Lewicki, H.J. Dyson, V.A. Feher, N. Assa-Munt and P.E. Wright (1991) *J. Virol.* **65**, 1727-1734.
126. Evidence for two interconverting protein isomers in the methotrexate complex of dihydrofolate reductase from *Escherichia coli*. C.J. Falzone, P.E. Wright and S.J. Benkovic (1991) *Biochemistry* **30**, 2184-2191.
127. Sensitivity improvement in proton detected two-dimensional heteronuclear correlation NMR spectroscopy. A.G. Palmer III, J. Cavanagh, P.E. Wright and M. Rance (1991) *J. Magn. Reson.* **93**, 151-170.
128. Intramolecular motions of a zinc finger DNA-binding domain from *Xfin* characterized by proton-detected natural abundance ¹³C heteronuclear NMR spectroscopy. A.G. Palmer III, M. Rance and P.E. Wright (1991) *J. Amer. Chem. Soc.* **113**, 4371-4380.
129. Polypeptide backbone resonance assignments and secondary structure of *Bacillus subtilis* enzyme III^{glc} determined by two-dimensional and three-dimensional heteronuclear NMR spectroscopy. W.J. Fairbrother, J. Cavanagh, H.J. Dyson, A.G. Palmer, S.L. Sutrina, J. Reizer, M.H. Saier, and P.E. Wright (1991) *Biochemistry* **30**, 6896-6907.
130. Measurement of relaxation rate constants for methyl groups by proton-detected heteronuclear NMR spectroscopy. A.G. Palmer III, P.E. Wright and M. Rance (1991) *Chem. Phys. Lett.* **185**, 41-46.
131. High-resolution solution structure of reduced French bean plastocyanin and comparison with the crystal structure of poplar plastocyanin. J.M. Moore, C. Lepre, G.P. Gippert, W.J. Chazin, D.A. Case and P.E. Wright (1991) *J. Mol. Biol.* **221**, 533-555.
132. Solution structures of DNA-binding domains of eukaryotic transcription factors. P.E. Wright (1992) In: *Transcriptional Regulation*, Eds. K. Yamamoto and S. McKnight, Cold Spring Harbor Laboratory Press, pp. 579-597.
133. Suppression of the effects of cross-correlation between dipolar and anisotropic chemical shift relaxation mechanisms in the measurement of spin-spin relaxation rates. A.G. Palmer III, N.J. Skelton, W.J. Chazin, P.E. Wright and M. Rance (1992) *Mol. Phys.*, **75**, 699-711.
134. A comparison of the requirements for pre-formed secondary structure in proteins with different structures in the folded state. H.J. Dyson and P.E. Wright (1992) In: *Proceedings of Seventh Conversation in Biomolecular Stereodynamics. Structure and Function, Volume 2: Proteins*, Eds. R.H. Sarma and M.H. Sarma, Adenine Press, Albany, NY, pp. 113-120.
135. Specific interactions of the first three zinc fingers of TFIIIA with the internal control region of the *Xenopus* 5S RNA gene. X. Liao, K. Clemens, L. Tennant, P.E. Wright and J. Gottesfeld (1992) *J. Mol. Biol.* **223**, 857-871.
136. Immunogenic peptides corresponding to the dominant antigenic region alanine-597 to cysteine-619 in the transmembrane protein of simian immunodeficiency virus have a propensity to fold in aqueous solution. H.J. Dyson, E. Norrby, K. Hoey, D.E. Parks, R.A. Lerner and P.E. Wright (1992) *Biochemistry* **31**, 1458-1463.
137. Improved resolution in three-dimensional constant-time triple resonance NMR spectroscopy of proteins. A.G. Palmer III, W.J. Fairbrother, J. Cavanagh, P.E. Wright and M. Rance (1992) *J. Biomol. NMR* **2**, 103-108.
138. Low resolution solution structure of the *Bacillus subtilis* glucose permease IIA domain derived from heteronuclear three-dimensional NMR spectroscopy. W.J. Fairbrother, G.P. Gippert, J. Reizer, M.H. Saier Jr. and P.E. Wright (1992) *FEBS Lett.* **296**, 148-152.

139. Structural determinants of Cys₂ His₂ zinc fingers. R.J. Mortishire-Smith, M.S. Lee, L. Bolinger and P.E. Wright (1992) *FEBS Lett.* **296**, 11-15.
140. Folding of peptide fragments comprising the complete sequence of proteins: Models for initiation of protein folding. I. Myohemerythrin. H.J. Dyson, G. Merutka, J.P. Waltho, R.A. Lerner and P.E. Wright (1992) *J. Mol. Biol.* **226**, 795-817.
141. Folding of peptide fragments comprising the complete sequence of proteins: Models for initiation of protein folding. II. Plastocyanin. H.J. Dyson, J.R. Sayre, H-C. Shin, G. Merutka, R.A. Lerner and P.E. Wright (1992) *J. Mol. Biol.* **226**, 819-835.
142. Relationship between ¹H and ¹³C NMR chemical shifts and the secondary and tertiary structure of a zinc finger peptide. M.S. Lee, A.G. Palmer III and P.E. Wright (1992) *J. Biomol. NMR* **2**, 307-322.
143. Backbone dynamics of the *Bacillus subtilis* glucose permease IIA domain determined from ¹⁵N NMR relaxation measurements. M.J. Stone, W.J. Fairbrother, A.G. Palmer III, J. Reizer, M.H. Saier Jr. and P.E. Wright (1992) *Biochemistry* **31**, 4394-4406.
144. Assignment of the aliphatic ¹H and ¹³C resonances of the *Bacillus subtilis* glucose permease IIA domain using double- and triple-resonance heteronuclear three-dimensional NMR spectroscopy. W.J. Fairbrother, A.G. Palmer, III, M. Rance, J. Reizer, M.H. Saier, Jr. and P.E. Wright (1992) *Biochemistry* **31**, 4413-4425.
145. Conformation and dynamics of an Fab'-bound peptide by isotope-edited NMR spectroscopy. P. Tsang, M. Rance, T.M. Fieser, J.M. Ostresh, R.A. Houghten, R.A. Lerner and P.E. Wright (1992) *Biochemistry* **31**, 3862-3871.
146. The zinc finger motif: Conservation of chemical shifts and correlation with structure. M.S. Lee, R.J. Mortishire-Smith, and P.E. Wright (1992) *FEBS Lett.* **309**, 29-32.
147. Functional role of a mobile loop of *Escherichia coli* dihydrofolate reductase in transition state stabilization. L. Li, C.J. Falzone, P.E. Wright and S.J. Benkovic (1992) *Biochemistry* **31**, 7826-7833.
148. Definition of the binding sites of individual zinc fingers in the TFIIIA-5S RNA gene complex. K.R. Clemens, X. Liao, V. Wolf, P.E. Wright and J.M. Gottesfeld (1992) *Proc. Natl Acad. Sci. USA* **89**, 10822-10826.
149. Determination of high resolution NMR structures of proteins. D.A. Case and P.E. Wright (1993) *In: NMR of Proteins*, Eds. G.M. Clore and A.M. Gronenborn, MacMillan, pp. 53-91.
150. Comparison of backbone and tryptophan sidechain dynamics of reduced and oxidized *E. coli* thioredoxin using ¹⁵N NMR relaxation measurements. M.J. Stone, K. Chandrasekhar, A. Holmgren, P.E. Wright and H.J. Dyson (1993) *Biochemistry* **32**, 426-435.
151. Three-dimensional structures of the central regulatory proteins of the bacterial phosphotransferase system, HPr and enzyme IIA^{glc}. Y. Chen, W.J. Fairbrother and P.E. Wright (1993) *J. Cell. Biochem.*, **51**, 75-82.
152. Mapping of the binding interfaces of the proteins of the bacterial phosphotransferase system, HPr and IIA^{glc}. Y. Chen, J. Reizer, M.H. Saier, W.J. Fairbrother and P.E. Wright (1993) *Biochemistry* **32**, 32-37.
153. Assignments of ¹H, ¹⁵N and ¹³C resonances, identification of elements of secondary structure and determination of the global fold of the DNA binding domain of GAL4. M. Shirakawa, W.J. Fairbrother, Y. Serikawa, T. Ohkubo, Y. Kyogoku and P.E. Wright (1993) *Biochemistry* **32**, 2144-2153.
154. Molecular basis for specific recognition of both RNA and DNA by a zinc finger protein. K.R. Clemens, V. Wolf, S.J. McBryant, P. Zhang, X. Liao, P.E. Wright and J.M. Gottesfeld (1993) *Science* **260**, 530-533.

155. The solution structure of the Oct-1 POU-specific domain reveals a striking similarity to the bacteriophage λ repressor DNA-binding domain. N. Assa-Munt, R. Mortishire-Smith, R. Aurora, W. Herr and P.E. Wright (1993) *Cell* **73**, 193-205.
156. Peptide conformation and protein folding. H.J. Dyson and P.E. Wright (1993) *Curr. Opin. Struct. Biol.* **3**, 60-65.
157. Structure of the retinoid X receptor α DNA-binding domain: A helix required for homodimeric DNA binding. M.S. Lee, S.A. Kliewer, J. Provencal, P.E. Wright and R.M. Evans (1993) *Science* **260**, 1117-1121.
158. Characterization of amino acid side chain dynamics in a zinc finger peptide using ^{13}C NMR spectroscopy and time-resolved fluorescence spectroscopy. A.G. Palmer, III, R.A. Hochstrasser, D.P. Millar, M. Rance and P.E. Wright (1993) *J. Amer. Chem. Soc.* **115**, 6333-6345.
159. Peptide models of protein folding initiation sites. 1. Secondary structure formation by peptides corresponding to the G- and H-helices of myoglobin. J.P. Waltho, V.A. Feher, G. Merutka, H.J. Dyson and P.E. Wright (1993) *Biochemistry* **32**, 6337-6347.
160. Peptide models of protein folding initiation sites. 2. The G-H turn region of myoglobin acts as a helix stop signal. H.-C. Shin, G. Merutka, J.P. Waltho, P.E. Wright and H. J. Dyson (1993) *Biochemistry* **32**, 6348-6355.
161. Peptide models of protein folding initiation sites. 3. The G-H helical hairpin of myoglobin. H.-C. Shin, G. Merutka, J.P. Waltho, L.L. Tennant, H.J. Dyson and P.E. Wright (1993) *Biochemistry* **32**, 6356-6364.
162. Determination of local ligand conformations in slowly tumbling proteins by homonuclear 2D and 3D NMR: Application to heme propionates in leghemoglobin. D. Morikis, R. Brüsweiler and P.E. Wright (1993) *J. Amer. Chem. Soc.* **115**, 6238-6246.
163. Electrostatic calculations of side-chain pK_a values in myoglobin and comparison with NMR data for histidines. D. Bashford, D.A. Case, C. Dalvit, L. Tennant and P.E. Wright (1993) *Biochemistry* **32**, 8045-8056.
164. Binding of a high-energy substrate conformer in antibody catalysis. A.P. Campbell, T.M. Tarasow, W. Massefski, P.E. Wright, and D. Hilvert (1993) *Proc. Natl Acad. Sci. USA* **90**, 8663-8667.
165. Formation of a molten globule intermediate in the kinetic folding pathway of apomyoglobin. P.A. Jennings and P.E. Wright (1993) *Science* **262**, 892-896.
166. Characterization of a folding intermediate of apoplastocyanin trapped by proline isomerization. S. Koide, H.J. Dyson and P.E. Wright (1993) *Biochemistry* **32**, 12299-12310.
167. NMR evidence for multiple conformations in a highly helical model peptide. G. Merutka, D. Morikis, R. Brüsweiler and P.E. Wright (1993) *Biochemistry* **32**, 13089-13097.
168. Biophysical methods: Editorial overview. J.L. Smith and P.E. Wright (1993) *Curr. Opin. Struct. Biol.* **3**, 723-724.
169. Dynamics of a flexible loop in dihydrofolate reductase from *Escherichia coli* and its implication for catalysis. C.J. Falzone, P.E. Wright and S.J. Benkovic (1994) *Biochemistry* **33**, 439-442.
170. POU domains and homeodomains. P.E. Wright (1994) *Curr. Opin. Struct. Biol.* **4**, 22-27.
171. ^1H resonance assignments and secondary structure of the carbon monoxide complex of soybean leghemoglobin determined by homonuclear 2D- and 3D-NMR spectroscopy. D. Morikis, C.A. Lepre and P.E. Wright (1994) *Eur. J. Biochem.* **219**, 611-626.
172. ^1H , ^{15}N , and ^{13}C resonance assignments, secondary structure, and the conformation of substrate in the binary folate complex of *Escherichia coli* dihydrofolate reductase. C.J. Falzone, J. Cavanagh, M. Cowart, A.G. Palmer III, S.J. Benkovic, C.R. Matthews and P.E. Wright (1994) *J. Biomol. NMR* **4**, 349-366.

173. ^1H , ^{15}N and ^{13}C resonance assignments for the first three zinc fingers of transcription factor IIIA. X. Liao, K.R. Clemens, J. Cavanagh, L. Tennant and P.E. Wright (1994) *J. Biomol. NMR* **4**, 433-454.
174. ^1H and ^{15}N resonance assignments and secondary structure of the carbon monoxide complex of sperm whale myoglobin. Y. Theriault, T.C. Pochapsky, C. Dalvit, M. Chiu, S.G. Sligar and P.E. Wright (1994) *J. Biomol. NMR*, **4**, 491-504.
175. Detecting nascent structures in solution using NMR. H.J. Dyson, J. Yao, and P.E. Wright (1994) *In: Proceedings of the 13th American Peptide Symposium. Peptides: Chemistry, Structure, and Biology* Eds. R.S. Hodges and J.A. Smith, ESCOM, Leiden, The Netherlands, 1093-1095.
176. The folding pathway of apomyoglobin. P.A. Jennings, H.J. Dyson and P.E. Wright (1994) *In: Statistical Mechanics, Protein Structure, and Protein Substrate Interactions*, NATO ARW, Ed. S. Doniach, Plenum Press, New York, 7-18.
177. Use of chemical shifts and coupling constants in NMR structural studies on peptides and proteins. D.A. Case, H.J. Dyson and P.E. Wright (1994) *In: Methods in Enzymology*, **239**, 392-416.
178. Protein structure calculation using NMR restraints. H.J. Dyson and P.E. Wright (1994) in *Two-dimensional NMR spectroscopy: Applications for chemists and biochemists*, Eds. W.R. Croasmun and R. Carlson, VCH, New York, 655-698.
179. NMR assignments and secondary structure of the retinoid X receptor \square DNA-binding domain: Evidence for the novel C-terminal helix. M.S. Lee, D.S. Sem, S.A. Kliewer, J. Provencal, R.M. Evans and P.E. Wright (1994) *Eur. J. Biochem.*, **224**, 639-650.
180. High-resolution solution structures of oxidized and reduced *Escherichia coli* thioredoxin. M-F. Jeng, A.P. Campbell, T. Begley, A. Holmgren, D.A. Case, P.E. Wright and H.J. Dyson (1994) *Struct.*, **2**, 853-868.
181. Solution structure of carbonmonoxy myoglobin determined from NMR distance and chemical shift constraints. K. Ösapay, Y. Theriault, P.E. Wright and D.A. Case (1994) *J. Mol. Biol.*, **244**, 183-197.
182. Relative contributions of the zinc fingers of transcription factor IIIA to the energetics of DNA binding. K.R. Clemens, P. Zhang, X. Liao, S.J. McBryant, P.E. Wright and J.M. Gottesfeld (1994) *J. Mol. Biol.*, **244**, 23-35.
183. NMR order parameters of biomolecules: A new analytical representation and application to the Gaussian axial fluctuation model. R. Brüschweiler and P.E. Wright (1994) *J. Am. Chem. Soc.*, **116**, 8426-8427.
184. Stabilization of a type VI turn in a family of linear peptides in water solution. J. Yao, V.A. Feher, F. Espejo, M.T. Reymond, P.E. Wright and H.J. Dyson (1994) *J. Mol. Biol.*, **243**, 736-753.
185. Three-dimensional structure of a type VI turn in a linear peptide in water solution: Evidence for stacking of aromatic rings as a major stabilizing factor. J. Yao, H.J. Dyson and P.E. Wright (1994) *J. Mol. Biol.*, **243**, 754-766.
186. Differential side chain hydration in a linear peptide containing a type VI turn. J. Yao, R. Brüschweiler, H.J. Dyson and P.E. Wright (1994) *J. Am. Chem. Soc.*, **116**, 12051-12052.
187. Water self-diffusion model for protein-water NMR cross relaxation. R. Brüschweiler and P.E. Wright (1994) *Chem. Phys. Lett.* **229**, 75-81.
188. Antigenic peptides. H.J. Dyson and P.E. Wright (1995) *FASEB J.*, **9**, 37-42.
189. 'Random coil' ^1H chemical shifts obtained as a function of temperature and trifluoroethanol concentration for the peptide series GGXGG. G. Merutka, H.J. Dyson and P.E. Wright, (1995) *J. Biomol. NMR*, **5**, 14-24.

190. Hydration of the partially folded peptide RN-24 studied by multi-dimensional NMR. R. Brüschweiler, D. Morikis and P.E. Wright (1995) *J Biomol. NMR*, **5**, 353-356.
191. Isotope labeling for NMR studies of macromolecular structure and interactions. P.E. Wright (1994) *Stable Isotope Applications in Biomolecular Structure and Mechanisms*, Eds. J. Trehwella, T.A. Cross, C.J. Unkefer, Los Alamos National Laboratory, Los Alamos, 1-11.
192. Interaction of the RNA binding fingers of *Xenopus* transcription factor IIIA with specific regions of 5S ribosomal RNA. S.J. McBryant, B. Gedulin, A. Leresche, M.P. Foster, P.E. Wright, P.J. Romaniuk and J.M. Gottesfeld (1995) *J. Mol. Biol.*, **248**, 44-57.
193. Recombinant soluble human tissue factor secreted by *Saccharomyces cerevisiae* and refolded from *E.coli* inclusion bodies: glycosylation of mutants, activity and physical characterization. M.J. Stone, W. Ruf, D.J. Miles, T.S. Edgington and P.E. Wright (1995) *Biochem. J.*, **310**, (Part 2), 605-614.
194. Long-range motional restrictions in a multidomain zinc-finger protein from anisotropic tumbling. R. Brüschweiler, X. Liao and P.E. Wright (1995) *Science* **268**, 886-889.
195. Measurement of intrinsic exchange rates of amide protons in a ¹⁵N-labeled peptide. S. Koide, W. Jahnke and P.E. Wright (1995) *J. Biomol. NMR*, **6**, 306-312.
196. Dynamics of the dihydrofolate reductase-folate complex: Catalytic sites and regions known to undergo conformational change exhibit diverse dynamical features. D.M. Epstein, S.J. Benkovic and P.E. Wright (1995) *Biochemistry*, **34**, 11037-11048.
197. The radius of gyration of an apomyoglobin folding intermediate. D. Eliezer, P.A. Jennings, P.E. Wright, S. Doniach, K.O. Hodgson and H. Tsuruta (1995) *Science* **270**, 487-488.
198. Structural basis for DNA bending by the architectural transcription factor LEF-1. J.J. Love, X. Li, D.A. Case, K. Giese, R. Grosschedl and P.E. Wright (1995) *Nature* **376**, 791-795.
199. Overexpression of myoglobin and assignment of its amide, C_α and C_β resonances. P.A. Jennings, M.J. Stone and P.E. Wright (1995) *J. Biomol. NMR* **6**, 271-276.
200. Proteins & Protein Fragments: Folding. H.J. Dyson and P.E. Wright (1996) **In:** *Encyclopedia of NMR*, (D.M. Grant and R.K. Harris, Eds.), Wiley, New York pp. 3811-3820.
201. NMR structural studies of flexible molecules. P.E. Wright and H.J. Dyson (1996) in *NMR as a Structural Tool for Macromolecules* (B.D. Nageswara Rao and M.D. Kemple, Eds) Plenum Publishing Corp., New York, pp. 245-249.
201. Hydrogen exchange in the carbon monoxide complex of soybean leghemoglobin. D. Morikis and P.E. Wright, (1996) *Eur. J. Biochem.* **237**, 212-220.
202. Probing protein/protein interactions with mass spectrometry and isotopic labeling: Analysis of the p21/Cdk2 complex. R.W. Kriwacki, J. Wu, G. Suizdak and P.E. Wright (1996) *J. Am. Chem. Soc.* **118**, 5320-5321.
203. Insights into protein folding from NMR. H.J. Dyson and P.E. Wright (1996) *Ann. Rev. Phys. Chem.* **47**, 369-395.
204. Assessment of major and minor groove DNA interactions by the zinc fingers of *Xenopus* transcription factor IIIA. S.J. McBryant, B. Gedulin, K.R. Clemens, P.E. Wright and J.M. Gottesfeld (1996) *Nucl. Acids Res.* **24**, 2567-2574.
205. Determination of the structures of proteins and protein complexes in solution using NMR. P.E. Wright, M. Foster, X. Li, J. Love and D. Wuttke (1996) *BioJapan '96, Symp. Proc.* 401-408.
206. Biophysical methods: Faster and bigger. P.E. Wright and J.L. Smith (1996) *Cur. Opin. Struct. Biol.* **6**, 583-584.

207. Structural studies of p21^{Waf1/Cip1/Sdi1} in the free and Cdk2-bound state: Conformational disorder mediates binding diversity. R.W. Kriwacki, L. Hengst, L. Tennant, S.I. Reed and P.E. Wright (1996) *Proc. Natl Acad. Sci. USA* **93**, 11504-11509.
208. Is apomyoglobin a molten globule? Structural characterization by NMR. D. Eliezer and P.E. Wright (1996) *J. Mol. Biol.* **263**, 531-538.
209. Gene synthesis, high level expression and assignment of backbone ¹⁵N and ¹³C resonances of soybean leghemoglobin. S. Prytulla, H.J. Dyson and P.E. Wright (1996) *FEBS Lett.* **399**, 283-289.
210. Folding propensities of peptide fragments of myoglobin. M.T. Reymond, G. Merutka, H.J. Dyson and P.E. Wright (1997) *Protein Sci.* **6**, 706-716.
211. Contribution of increased length and intact capping sequences to the conformational preference for helix in a 31-residue peptide from the C-terminus of myohemerythrin. M.T. Reymond, S. Huo, B. Duggan, P.E. Wright and H.J. Dyson (1997) *Biochemistry* **36**, 5234-5244.
212. Absence of a stable intermediate on the folding pathway of protein A. Y. Bai, A. Karimi, H.J. Dyson and P.E. Wright (1997) *Protein Sci.* **6**, 1449-1457.
213. NMR spectroscopic studies of the DNA-binding domain of the monomer-binding nuclear orphan receptor, human estrogen related receptor-2. The carboxy-terminal extension to the zinc-finger region is unstructured in the free form of the protein. D.S. Sem, D.R. Casimiro, S.A. Kliewer, J. Provencal, R.M. Evans and P.E. Wright (1997) *J. Biol. Chem.* **272**, 18038-18043.
214. Domain packing and dynamics in the DNA complex of the N-terminal zinc fingers of TFIIIA. M.P. Foster, D.S. Wuttke, I. Radhakrishnan, D.A. Case, J.M. Gottesfeld and P.E. Wright (1997) *Nat. Struct. Biol.* **4**, 605-608.
215. Solution structure of the first three zinc fingers of TFIIIA bound to the cognate DNA sequence: Determinants of affinity and sequence specificity. D.S. Wuttke, M.P. Foster, D.A. Case, J.M. Gottesfeld and P.E. Wright (1997) *J. Mol. Biol.* **273**, 183-206.
216. Populating the equilibrium molten globule state of apomyoglobin under conditions suitable for structural characterization by NMR. D. Eliezer, P.A. Jennings, H.J. Dyson and P.E. Wright (1997) *FEBS Lett.* **417**, 92-96.
217. Probing protein structure using biochemical and biophysical methods: Proteolysis, matrix-assisted laser desorption/ionization mass spectrometry, high-performance liquid chromatography and size-exclusion chromatography of p21^{Waf1/Cip1/Sdi1}. R.W. Kriwacki, J. Wu, L. Tennant, P.E. Wright and G. Siuzdak (1997) *J. Chrom. A* **777**, 23-30.
218. PCR-based gene synthesis and protein NMR spectroscopy. D.R. Casimiro, P.E. Wright and H.J. Dyson (1997) *Struct.* **5**, 1407-1412.
219. Structure of the recombinant full-length hamster prion protein PrP (29-231): The N-terminus is highly flexible. D.G. Donne, J.H. Viles, D. Groth, I. Melhorn, T.L. James, F.E. Cohen, S.B. Prusiner, P.E. Wright and H.J. Dyson (1997) *Proc. Natl Acad. Sci., USA* **94**, 13452-13457.
220. Solution structure of the KIX domain of CBP bound to the transactivation domain of CREB: A model for activator:coactivator interactions. I. Radhakrishnan, G.C. Perez-Alvarado, H.J. Dyson, D. Parker, M.R. Montminy and P.E. Wright (1997) *Cell* **91**, 741-752.
221. Chemical shift dispersion and secondary structure prediction in unfolded and partly-folded proteins. J. Yao, H.J. Dyson and P.E. Wright (1997) *FEBS Lett.* **419**, 285-289.

222. Structural and dynamic characterization of partially folded states of apomyoglobin: Implications for protein folding. D. Eliezer, J. Yao, H.J. Dyson and P.E. Wright (1998) *Nat. Struct. Biol.* **5**, 148-155.
223. A NOESY-HSQC simulation program, SPIRIT. L. Zhu, H.J. Dyson and P.E. Wright (1998) *J. Biomol. NMR* **11**, 17-29.
224. High-resolution solution structure of *Bacillus subtilis* IIA^{glc}. Y. Chen, D.A. Case, J. Reizer, M.H. Saier, Jr. and P.E. Wright (1998) *Proteins: Structure, Function and Genetics* **31**, 258-270.
225. Recommendations for the presentation of NMR structures of proteins and nucleic acids. J.L. Markley, A. Bax, Y. Arata, C.W. Hilbers, R. Kaptein, B.D. Sykes, P.E. Wright and K. Wüthrich (1998) *Pure Appl. Chem.* **70**, 117-142.
226. Conformational preferences in the Ser133-phosphorylated and non-phosphorylated forms of the kinase inducible transactivation domain of CREB. I. Radhakrishnan, G.C. Perez-Alvarado, H.J. Dyson and P.E. Wright (1998) *FEBS Lett.* **430**, 317-322.
227. Equilibrium NMR studies of unfolded and partially folded proteins. H.J. Dyson and P.E. Wright (1998) *Nat. Struct. Biol.* **5**, 499-503.
228. Distributed torsion angle grid search in high dimensions: A systematic approach to NMR structure determination. G.P. Gippert, P.E. Wright and D.A. Case (1998) *J. Biomol. NMR* **11**, 241-263.
229. High resolution solution structure of the retinoid X receptor DNA binding domain. S.M.A. Holmbeck, M.P. Foster, D.R. Casimiro, D. Sem, H.J. Dyson and P.E. Wright (1998) *J. Mol. Biol.* **281**, 271-284.
230. Chemical shift as a probe of molecular interfaces: NMR studies of DNA binding by three amino-terminal zinc finger domains from transcription factor IIIA. M.P. Foster, D.S. Wuttke, K.R. Clemens, W. Jahnke, I. Radhakrishnan, L. Tennant, M. Reymond, J. Chung and P.E. Wright (1998) *J. Biomol. NMR* **12**, 51-71.
231. ¹H, ¹³C and ¹⁵N NMR backbone assignments of 25.5 kDa metallo-β-lactamase from bacteroides fragilis. S.D.B. Scrofani, P.E. Wright and H.J. Dyson (1998) *J. Biomol. NMR* **12**, 201-202.
232. Sequence requirements for stabilization of a peptide reverse turn in water solution: proline is not essential for stability. H.J. Dyson, L. Bolinger, V.A. Feher, J.J. Osterhout, Jr., J. Yao and P.E. Wright (1998) *Eur. J. Biochem.* **255**, 462-471.
233. Analysis of an activator:Coactivator complex reveals an essential role for secondary structure in transcriptional activation. D. Parker, U. Jhala, I. Radhakrishnan, M.B. Yaffe, C. Reyes, A.I. Shulman, L.C. Cantley, P.E. Wright and M. Montminy (1998). *Mol. Cell* **2**, 353-359.
234. The identification of metal-binding ligand residues in metalloproteins using nuclear magnetic resonance spectroscopy. S.D.B. Scrofani, P.E. Wright and H.J. Dyson (1998) *Protein Sci.* **7**, 2476-2479.
235. DNA-induced conformational changes are the basis for cooperative dimerization by the DNA binding domain of the retinoid-X-receptor. S.M.A. Holmbeck, H.J. Dyson and P.E. Wright (1998) *J. Mol. Biol.* **284**, 533-539.
236. Effect of H helix destabilizing mutations on the kinetic and equilibrium folding of apomyoglobin. S. Cavagnero, H.J. Dyson, P.E. Wright (1999) *J. Mol. Biol.* **285**, 269-282.
237. Quench-flow experiments combined with mass spectrometry show apomyoglobin folds through an obligatory intermediate. V. Tsui, C. Garcia, S. Cavagnero, G. Siuzdak, H.J. Dyson and P.E. Wright (1999) *Protein Sci.* **8**, 45-49.
238. Copper binding to the prion protein: Structural implications of four identical cooperative binding sites. J.H. Viles, F.E. Cohen, S.B. Prusiner, D.B. Goodin, P.E. Wright and H.J. Dyson (1999) *Proc. Natl Acad. Sci., USA* **96**, 2042-2047.

239. Structural analyses of CREB-CBP transcriptional activator-coactivator complexes by NMR spectroscopy: Implications for mapping the boundaries of structural domains. I. Radhakrishnan, G.C. Perez-Alvarado, D. Parker, H.J. Dyson, M. Montminy and P.E. Wright (1999) *J. Mol. Biol.* **287**, 859-865.
240. Improved low pH bicelle system for orienting macromolecules over a wide temperature range. S. Cavagnero, H.J. Dyson and P.E. Wright (1999) *J. Biomol. NMR* **13**, 387-391.
241. Association between the first two immunoglobulin-like domains of the neural cell adhesion molecule N-CAM. A.R. Atkins, M.J. Osborne, H.A. Lashuel, G.M. Edelman, P.E. Wright, B.A. Cunningham and H.J. Dyson (1999) *FEBS Lett.* **451**, 162-168.
242. Role of secondary structure in discrimination between constitutive and inducible activators. D. Parker, M. Rivera, T. Zor, A. Henrion-Caude, I. Radhakrishnan, A. Kumar, L.H. Shapiro, P.E. Wright, M. Montminy and P.K. Brindle (1999) *Mol. Cell. Biol.* **19**, 5601-5607.
243. Identification of a minimal domain of 5S ribosomal RNA sufficient for high affinity interactions with the RNA-specific zinc fingers of transcription factor IIIA. L.S. Neely, B.M. Lee, J. Xu, P.E. Wright and J.M. Gottesfeld (1999) *J. Mol. Biol.* **291**, 549-560.
244. Backbone resonance assignments for the Fv fragment of the catalytic antibody NPN43C9 with bound *p*-nitrophenol. G.J.A. Kroon, M.A. Martinez-Yamout, J.F. Krebs, J. Chung, H.J. Dyson and P.E. Wright (1999) *J. Biomol. NMR* **15**, 83-84.
245. Assignment of ^1H , ^{13}C , and ^{15}N resonances of reduced *Escherichia coli* glutaredoxin 2. B. Xia, J. Chung, A. Vlamis-Gardikas, A. Holmgren, P.E. Wright and H.J. Dyson (1999) *J. Biomol. NMR* **14**, 197-198.
246. Characterization of monomeric and dimeric B domain of staphylococcal protein A: Sources of stabilization of a 3-helix bundle protein. A. Karimi, M. Matsumura, P.E. Wright and H.J. Dyson (1999) *J. Pept. Res.* **54**, 344-352.
247. Intrinsically unstructured proteins: re-assessing the protein structure-function paradigm. P.E. Wright and H.J. Dyson (1999) *J. Mol. Biol.* **293**, 321-331.
248. Inherent flexibility in a potent inhibitor of blood coagulation, recombinant nematode anticoagulant protein c2. B.M. Duggan, H.J. Dyson and P.E. Wright (1999) *Eur. J. Biochem.* **265**, 539-548.
249. NMR characterization of the metallo- β -lactamase from *bacteroides fragilis* and its interaction with a tight-binding inhibitor: Role of an active site loop. S.D.B. Scrofani, J. Chung, J.J.A. Huntley, S.J. Benkovic, P.E. Wright and H.J. Dyson (1999) *Biochemistry*, **38**, 14507-14514.
250. Determination of Fe-ligand bond lengths and the Fe-N-O bond angles in soybean ferrous and ferric nitrosylhemoglobin α using multiple-scattering XAFS analyses. A.M. Rich, P.J. Ellis, L. Tennant, P.E. Wright, R.S. Armstrong and P.A. Lay (1999) *Biochemistry*, **38**, 16491-16499.
251. NMR solution structure of the inserted domain of human leukocyte function associated antigen-1. G.B. Legge, R.W. Kriwacki, J. Chung, U. Hommel, P. Ramage, D.A. Case, H.J. Dyson and P.E. Wright (2000) *J. Mol. Biol.*, **295**, 1251-1264.
252. DNA-induced α -helix capping in conserved linker sequences is a determinant of binding affinity in Cys₂-His₂ zinc fingers. J.H. Laity, H.J. Dyson and P.E. Wright (2000) *J. Mol. Biol.*, **295**, 719-727.
253. Amide proton hydrogen exchange rates for sperm whale myoglobin obtained from ^{15}N - ^1H NMR spectra. S. Cavagnero, Y. Thériault, S. Narula, H.J. Dyson and P.E. Wright (2000) *Protein Sci.*, **9**, 186-193.
254. Assessment of zinc finger orientations by residual dipolar coupling constants. V. Tsui, L. Zhu, T-H. Huang, P.E. Wright and D.A. Case (2000) *J. Biomol. NMR*, **16**, 9-21.

255. Native and non-native secondary structure and dynamics in the pH 4 intermediate of apomyoglobin. D. Eliezer, J. Chung, H.J. Dyson and P.E. Wright (2000) *Biochemistry*, **39**, 2894-2901.
256. Identification of the regions involved in DNA-binding by the mouse PEBP2 α protein. G.C. Perez-Alvarado, A. Munnerlyn, H.J. Dyson, R. Grosschedl and P.E. Wright (2000) *FEBS Letters*, **470**, 125-130.
257. Assignment of ^1H , ^{13}C and ^{15}N resonances of the I-domain of human leukocyte function associated antigen-1. R.W. Kriwacki, G.B. Legge, U. Hommel, P. Ramage, J. Chung, L.L. Tennant, P.E. Wright and H.J. Dyson (2000) *J. Biomol. NMR*, **16**, 271-272.
258. Alternative splicing of Wilms' tumor suppressor protein modulates DNA binding activity through isoform-specific DNA-induced conformational changes. J.H. Laity, J. Chung, H.J. Dyson and P.E. Wright (2000) *Biochemistry*, **39**, 5341-5348.
259. Backbone H^{N} , N , C^{α} , C' and C^{β} assignments of the 19 kDa DHFR/NADPH complex at 9°C and pH 7.6. E. Zaborowski, J. Chung, G. Kroon, H.J. Dyson and P.E. Wright (2000) *J. Biomol. NMR*, **16**, 349-350.
260. Solution structure of the cysteine-rich domain of the *Escherichia coli* chaperone protein DnaJ. M. Martinez-Yamout, G.B. Legge, O. Zhang, P.E. Wright and H.J. Dyson (2000) *J. Mol. Biol.*, **300**, 805-818.
261. Conservation of folding pathways in evolutionarily distant globin sequences. C. Nishimura, S. Prytulla, H.J. Dyson and P.E. Wright (2000) *Nat. Struct. Biol.*, **7**, 679-686.
262. NMR and molecular dynamics studies of the hydration of a zinc finger-DNA complex. V. Tsui, I. Radhakrishnan, P.E. Wright and D.A. Case (2000) *J. Mol. Biol.*, **302**, 1101-1117.
263. Solution structure of the TAZ2 (CH3) domain of the transcriptional adaptor protein CBP. R.N. De Guzman, H.Y. Liu, M. Martinez-Yamout, H.J. Dyson and P.E. Wright (2000) *J. Mol. Biol.*, **303**, 243-253.
264. Changes in the apomyoglobin folding pathway caused by mutation of the distal histidine residue. C. Garcia, C. Nishimura, S. Cavagnero, H.J. Dyson and P.E. Wright (2000) *Biochemistry*, **39**, 11227-11237.
265. Solution structure and acetyl-lysine binding activity of the GCN5 bromodomain. B.P. Hudson, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2000) *J. Mol. Biol.*, **304**, 355-370.
266. Random coil chemical shifts in acidic 8 M urea: Implementation of random coil shift data in NMRview. S. Schwarzinger, G.J.A. Kroon, T.R. Foss, P.E. Wright, H.J. Dyson (2000) *J. Biomol. NMR*, **18**, 43-48.
267. Molecular basis for modulation of biological function by alternate splicing of the Wilms' tumor suppressor protein. J.H. Laity, H.J. Dyson and P.E. Wright (2000) *Proc. Natl Acad. Sci, USA*, **97**, 11932-11935.
268. Structure of the PHD zinc finger from human Williams-Beuren Syndrome transcription factor. J. Pascual, M. Martinez-Yamout, H.J. Dyson and P.E. Wright (2000) *J. Mol. Biol.*, **304**, 723-729.
269. Dynamics of the metallo- β -lactamase from *Bacteroides fragilis* in the presence and absence of a tight-binding inhibitor. J.J.A. Huntley, S.D.B. Scrofani, M.J. Osborne, P.E. Wright and H.J. Dyson (2000) *Biochemistry*, **39**, 13356-13364.
270. The folding process of apomyoglobin. P.E. Wright and R.L. Baldwin (2000) *In: Frontiers in Molecular Biology: Mechanisms of Protein Folding*, 2nd edition (R. Pain, ed.), Oxford University Press, UK pp. 309-329.
271. Zinc finger proteins: New insights into structural and functional diversity. J.H. Laity, B.M. Lee and P.E. Wright (2001) *Curr. Opin. Struct. Biol.*, **11**, 39-46.
272. Local structural plasticity of the prion protein. Analysis of NMR relaxation dynamics. J.H. Viles, D. Donne, G. Kroon, S.B. Prusiner, F.E. Cohen, H.J. Dyson and P.E. Wright (2001) *Biochemistry*, **40**, 2743-2753.

273. Two different neurodegenerative diseases caused by proteins with similar structures. H. Mo, R.C. Moore, F.E. Cohen, D. Westaway, S.B. Prusiner, P.E. Wright and H.J. Dyson (2001) *Proc. Natl Acad. Sci, USA*, **98**, 2352-2357.
274. Anisotropic rotational diffusion in model-free analysis for a ternary-DHFR complex. M.J. Osborne and P.E. Wright (2001) *J. Biomol. NMR*, **19**, 209-230.
275. Sequence-dependent correction of random coil NMR chemical shifts. S. Schwarzingler, G.J.A. Kroon, T.R. Foss, J. Chung, P.E. Wright and H.J. Dyson (2001) *J. Amer. Chem. Soc.*, **123**, 2970-2978.
276. SANE (structure assisted NOE evaluation): An automated model-based approach for NOE assignment. B.M. Duggan, G.B. Legge, H.J. Dyson and P.E. Wright (2001) *J. Biomol. NMR*, **19**, 321-329.
277. NMR structural and dynamic characterization of the acid-unfolded state of apomyoglobin provides insights into the early events in protein folding. J. Yao, J. Chung, D. Eliezer, P.E. Wright and H.J. Dyson (2001) *Biochemistry*, **40**, 3561-3571.
278. Structural and dynamic characterization of an unfolded state of poplar apo-plastocyanin formed under non-denaturing conditions. Y. Bai, J. Chung, H.J. Dyson and P.E. Wright (2001) *Protein Science*, **10**, 1056-1066.
279. Nuclear magnetic resonance methods for the elucidation of structure and dynamics in disordered states. H.J. Dyson and P.E. Wright (2001) *In: Methods in Enzymology*, (T.L. James, V. Dötsch, and U. Schmitz, eds.), Academic Press, San Diego **339**, 258-270.
280. Solution structure of *Escherichia coli* glutaredoxin-2 shows similarity to mammalian glutathione-S-transferases. B. Xia, A. Vlamis-Gardikas, A. Holmgren, P.E. Wright and H.J. Dyson (2001) *J. Mol. Biol.*, **310**, 907-918.
281. Solution structure of the third immunoglobulin domain of the neural cell adhesion molecule N-CAM: Can solution studies define the mechanism of homophilic binding? A.R. Atkins, J. Chung, S-P. Deechongkit, E.B. Little, G.M. Edelman, P.E. Wright, B.A. Cunningham and H.J. Dyson (2001) *J. Mol. Biol.*, **311**, 161-172.
282. Backbone dynamics in dihydrofolate reductase complexes: Role of loop flexibility in the catalytic mechanism. M.J. Osborne, J. Schnell, S.J. Benkovic, H.J. Dyson and P.E. Wright (2001) *Biochemistry*, **40**, 9846-9859. (*Faculty of 1000 Recommended paper, October 2001*).
283. Potential bias in NMR relaxation data introduced by peak intensity analysis and curve fitting methods. J.H. Viles, B.M. Duggan, E. Zaborowski, S. Schwarzingler, J.J.A. Huntley, G.J.A. Kroon, H.J. Dyson and P.E. Wright (2001) *J. Biomol. NMR*, **21**, 1-9.
284. Conformational and dynamic characterization of the molten globule state of an apomyoglobin mutant with an altered folding pathway. S. Cavagnero, C. Nishimura, S. Schwarzingler, H.J. Dyson and P.E. Wright (2001) *Biochemistry*, **40**, 14459-14467.
285. Mutual synergistic folding in recruitment of CBP/p300 by p160 nuclear receptor coactivators. S.J. Demarest, M. Martinez-Yamout, J. Chung, H. Chen, W. Xu, H.J. Dyson, R.M. Evans and P.E. Wright (2002) *Nature*, **415**, 549-553.
286. Coupling of folding and binding for unstructured proteins. H.J. Dyson and P.E. Wright (2002) *Curr. Opin. Struct. Biol.*, **12**, 54-60.
287. Solution NMR spectroscopy of [α - ^{15}N]lysine-labeled rhodopsin: The single peak observed in both conventional and TROSY-type HSQC spectra is ascribed to Lys-339 in the carboxyl-terminal peptide sequence. J. Klein-Seetharaman, P.J. Reeves, M.C. Loewen, E.V. Getmanova, J. Chung, H. Schwalbe, P.E. Wright and H.G. Khorana (2002) *Proc. Natl Acad. Sci, USA*, **99**, 3452-3457.

288. Structural basis for Hif-1 α /CBP recognition in the cellular hypoxic response. S.A. Dames, M. Martinez-Yamout, R.N. De Guzman, H.J. Dyson and P.E. Wright (2002) *Proc. Natl Acad. Sci, USA*, **99**, 5271-5276. (*Faculty of 1000 Must Read paper, May 2002*).
289. Assignment of a 15 kDa protein complex formed between the p160 coactivator ACTR and CREB binding protein. S.J. Demarest, J. Chung, H.J. Dyson and P.E. Wright (2002) *J. Biomol. NMR*, **22**, 377-378.
290. Comparison of protein solution structures refined by molecular dynamics simulation in vacuum, with a generalized Born model, and with explicit water. B. Xia, V. Tsui, D.A. Case, H.J. Dyson and P.E. Wright (2002), *J. Biomol. NMR*, **22**, 317-331.
291. Structure and dynamics of disordered proteins. H.J. Dyson and P.E. Wright (2002) *In: Encyclopedia of NMR*, (D.M. Grant & R.K. Harris, eds.), John Wiley & Sons, Ltd., **9**, 449-457.
292. High pressure NMR reveals that apomyoglobin is an equilibrium mixture from the native to the unfolded. R. Kitahara, H. Yamada, K. Akasaka and P.E. Wright (2002) *J. Mol. Biol.*, **320**, 311-319.
293. Mapping long-range contacts in a highly unfolded protein. M.A. Lietzow, M. Jamin, H.J. Dyson and P.E. Wright (2002) *J. Mol. Biol.*, **322**, 655-662.
294. Molecular hinges in protein folding: the urea-denatured state of apomyoglobin. S. Schwarzingler, P.E. Wright and H.J. Dyson (2002) *Biochemistry*, **41**, 12681-12686.
295. The apomyoglobin folding pathway revisited: Structural heterogeneity in the kinetic burst phase intermediate. C. Nishimura, H.J. Dyson and P.E. Wright (2002) *J. Mol. Biol.*, **322**, 483-489.
296. Roles of phosphorylation and helix propensity in the binding of the KIX domain of CREB-binding protein by constitutive (c-Myb) and inducible (CREB) activators. T. Zor, B.M. Mayr, H.J. Dyson, M.R. Montminy and P.E. Wright (2002) *J. Biol. Chem.*, **277**, 42241-42248.
297. Insights into the structure and dynamics of unfolded proteins from nuclear magnetic resonance. H. Jane Dyson and P.E. Wright (2002) *Adv. in Protein Chem.*, **62**, 311-40.
298. Cooperativity in transcription factor binding to the coactivator CREB-binding Protein (CBP). N.K. Goto, T. Zor, M. Martinez-Yamout, H.J. Dyson and P.E. Wright (2002) *J. Biol. Chem.*, **277**, 43168-43174.
299. Monomeric complex of human orphan estrogen related receptor-2 with DNA: A pseudo-dimer interface mediates extended half-site recognition. M.D. Gearhart, S.M.A. Holmbeck, R.M. Evans, H.J. Dyson and P.E. Wright (2003) *J. Mol. Biol.*, **327**, 819-832.
300. Folding of a β -sheet protein monitored by real-time NMR spectroscopy. M. Mizuguchi, G.J. Kroon, P.E. Wright and H.J. Dyson (2003) *J. Mol. Biol.*, **328**, 1161-1171.
301. Structure of the nuclear factor ALY: Insights into post-transcriptional regulatory and mRNA nuclear export processes. G.C. Perez-Alvarado, M. Martinez-Yamout, M.M. Allen, R. Grosschedl, H.J. Dyson and P.E. Wright (2003) *Biochemistry*, **42**, 7348-7357.
302. Role of a solvent-exposed tryptophan in the recognition and binding of antibiotic substrates for a metallo- β -lactamase. J.J.A. Huntley, W. Fast, S.J. Benkovic, P.E. Wright and H.J. Dyson (2003) *Protein Science*, **12**, 1368-1375.
303. Changes in structure and dynamics of the Fv fragment of a catalytic antibody upon binding of inhibitor. G.J.A. Kroon, H. Mo, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2003) *Protein Science*, **12**, 1386-1394.

304. Diagnostic chemical shift markers for loop conformation and substrate and cofactor binding in dihydrofolate reductase complexes. M.J. Osborne, R.P. Venkitakrishnan, H.J. Dyson and P.E. Wright (2003) *Protein Science*, **12**, 2230-2238.
305. Role of the B helix in early folding events in apomyoglobin: Evidence from site-directed mutagenesis for native-like long range interactions. C. Nishimura, P.E. Wright and H.J. Dyson (2003) *J. Mol. Biol.*, **334**, 293-307.
306. Packing, specificity and mutability at the binding interface between the p160 coactivator and CREB-binding protein. S.J. Demarest, S. Deechongkit, H.J. Dyson, R.M. Evans and P.E. Wright (2004) *Protein Science*, **13**, 203-210.
307. Effect of cofactor binding and loop conformation on side chain methyl dynamics in dihydrofolate reductase. J.R. Schnell, H.J. Dyson and P.E. Wright (2004) *Biochemistry*, **43**, 374-383.
308. Interaction of the TAZ1 domain of the CREB-binding protein with the activation domain of CITED2: Regulation by competition between intrinsically unstructured ligands for non-identical binding sites. R.N. De Guzman, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2004) *J. Biol. Chem.*, **279**, 3042-3049.
309. Recognition of the mRNA AU-rich element by the zinc finger domain of TIS11d. B.P. Hudson, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2004) *Nat. Struct. Biol.*, **11**, 257-264.
310. Solution structure of the KIX domain of CBP bound to the transactivation domain of c-Myb. T. Zor, R.N. De Guzman, H.J. Dyson and P.E. Wright (2004) *J. Mol. Biol.*, **337**, 521-534.
311. Structure, dynamics and catalytic function of dihydrofolate reductase. J.R. Schnell, H.J. Dyson and P.E. Wright (2004) *Ann. Rev. Biophys. Biomol. Struct.*, **33**, 119-140.
312. Model-free analysis of protein dynamics: assessment of accuracy and model selection protocols based on molecular dynamics simulation. J. Chen, C.L. Brooks, III and P.E. Wright (2004) *J. Biomolecular NMR*, **29**, 243-257.
313. The LEF-1 high-mobility group domain undergoes a disorder-to-order transition upon formation of a complex with cognate DNA. J.J. Love, X. Li, J. Chung, H.J. Dyson and P.E. Wright (2004) *Biochemistry*, **43**, 8725-8734.
314. The CBP/p300 TAZ1 domain in its native state is not a binding partner of MDM2. T. Matt, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2004) *Biochem. J.*, **381**, 685-691. (*Commentary: Biochem. J.* **381**, e3-e4, 2004; *Faculty of 1000 Must Read paper, August 2004*).
315. Structural characterization of unfolded states of apomyoglobin using residual dipolar couplings. R. Mohana-Borges, N.K. Goto, G.J.A. Kroon, H.J. Dyson and P.E. Wright (2004) *J. Mol. Biol.*, **340**, 1131-1142.
316. *Thematic issue: Biological Nuclear Magnetic Resonance*. H.J. Dyson and P.E. Wright, Guest Editors (2004) *Chemical Reviews* **104**, pp. 3517-3704.
317. Unfolded proteins and protein folding studied by NMR. H.J. Dyson and P.E. Wright (2004) *Chemical Reviews*, **104**, 3607-3622.
318. Conformational changes in the active site loops of dihydrofolate reductase during the catalytic cycle. R.P. Venkitakrishnan, E. Zaborowski, D. McElheny, S.J. Benkovic, H.J. Dyson and P.E. Wright (2004), *Biochemistry*, **43**, 16046-16055.
319. Structure and function of the CBP/p300 TAZ domains. R.N. De Guzman, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2004) *In: Zinc Finger Proteins: From Atomic Contact to Cellular Function*, Eds. S. Iuchi, N. Kuldell, Landes Bioscience, Georgetown, TX **17**, 114-122.

320. ZZ domain of CBP: an unusual zinc finger fold in a protein interaction module. G.B. Legge, M.A. Martinez-Yamout, D.M. Hambly, T. Trinh, B.M. Lee, H.J. Dyson and P.E. Wright (2004), *J. Mol. Biol.* **343**, 1081-1093.
321. CBP/p300 TAZ1 domain forms a structural scaffold for ligand binding. R.N. De Guzman, J.M. Wojciak, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2005), *Biochemistry* **44**, 490-497.
322. Intrinsically unstructured proteins and their function. H.J. Dyson and P.E. Wright (2005) *Nature Rev. Mol. Cell Biol.*, **6**, 197-208. (Highlighted in ESI Special Topics, February 2006, as a Fast Breaking Paper. <http://esi-topics.com>).
323. Inhibition of DNA binding by human estrogen related receptor-2 and estrogen receptor α with minor groove-binding polyamides. M.D. Gearhart, L. Dickinson, J. Ehley, C. Melander, P.B. Dervan, P.E. Wright and J.M. Gottesfeld (2005), *Biochemistry* **44**, 4196-4203.
324. Enhanced picture of protein-folding intermediates using organic solvents in H/D exchange and quench-flow experiments. C. Nishimura, H.J. Dyson and P.E. Wright (2005), *Proc. Natl Acad. Sci., USA* **102**, 4765-4770.
325. Defining the role of active-site loop fluctuations in dihydrofolate reductase catalysis. D. McElheny, J.R. Schnell, J.C. Lansing, H.J. Dyson and P.E. Wright (2005), *Proc. Natl Acad. Sci., USA* **102**, 5032-5037. (Faculty of 1000 Must Read paper, May 2005).
326. Elucidation of the protein folding landscape by NMR. H.J. Dyson and P.E. Wright (2005) *In: Methods in Enzymology*, (Thomas L. James, Ed.) Elsevier, Inc. **394**, 299-321.
327. Sequence determinants of a protein folding pathway. C. Nishimura, M.A. Lietzow, H.J. Dyson and P.E. Wright (2005), *J. Mol. Biol.* **351**, 383-392.
328. Solution structure of the N-terminal zinc fingers of the xenopus laevis double-stranded RNA-binding protein ZFa. H.M. Möller, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2005), *J. Mol. Biol.* **351**, 718-730.
329. Identification of native and non-native structure in kinetic folding intermediates of apomyoglobin. C. Nishimura, H.J. Dyson and P.E. Wright (2006), *J. Mol. Biol.* **355**, 139-156.
330. Structure of the *Escherichia coli* quorum sensing protein SdiA: Activation of the folding switch by acyl homoserine lactones. Y. Yao, M.A. Martinez-Yamout, T.J. Dickerson, A.P. Brogan, P.E. Wright and H.J. Dyson (2006), *J. Mol. Biol.* **355**, 262-273.
331. Structural basis for cooperative transcription factor binding to the CBP coactivator. R.N. De Guzman, N.K. Goto, H.J. Dyson and P.E. Wright (2006), *J. Mol. Biol.* **355**, 1005-1013.
332. NMR solution structure of the peptide fragment 1-30, derived from unprocessed mouse Doppel protein, in DHPC micelles. E. Papadopoulos, K. Oglecka, L. Mäler, J. Jarver, P.E. Wright, H.J. Dyson, and A. Gräslund (2006), *Biochemistry* **45**, 159-166.
333. Induced fit and lock and key recognition of 5S RNA by zinc fingers of transcription factor IIIA. B.M. Lee, J. Xu, B.K. Clarkson, M.A. Martinez-Yamout, H.J. Dyson, D.A. Case, J.M. Gottesfeld and P.E. Wright (2006), *J. Mol. Biol.* **357**, 275-291.
334. Is there an answer? According to current textbooks, a well-defined three-dimensional structure is a prerequisite for the function of a protein. Is this correct? H.J. Dyson and P.E. Wright (2006), *IUBMB Life* **58**, 107-109.
335. Localization of sites of interaction between p23 and Hsp90 in solution. M.A. Martinez-Yamout, R.P. Venkitakrishnan, N.E. Preece, G. Kroon, P.E. Wright and H.J. Dyson (2006), *J. Biol. Chem.* **281**, 14457-14464.
336. The role of hydrophobic interactions in initiation and propagation of protein folding. H.J. Dyson, P.E. Wright and H.A. Scheraga (2006), *Proc. Natl Acad. Sci., USA* **103**:13057-13061.

337. The dynamic energy landscape of dihydrofolate reductase catalysis. D.D. Boehr, D. McElheny, H.J. Dyson and P.E. Wright (2006), *Science* **313**, 1638-1642. (*Perspective by M. Vendruscolo and C.M. Dobson, Science 313, 1586, 2006; Faculty of 1000 Exceptional Paper*).
338. Solution structure of the Hdm2 C2H2C4 RING, a domain critical for ubiquitination of p53. M. Kostic, T. Matt, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2006) *J. Mol. Biol.* **363**, 433-450.
339. An NMR Perspective on Enzyme Dynamics. D.D. Boehr, H.J. Dyson and P.E. Wright (2006), *Chemical Reviews* **106**, 3055-3079.
340. PRAK is essential for *ras*-induced senescence and tumor suppression. P. Sun, N. Yoshizuka, L. New, B.A. Moser, Y. Li, R. Liao, C. Xie, J. Chen, Q. Deng, M. Martinez-Yamout, M-Q. Dong, C.G. Frangou, J.R. Yates, P.E. Wright and J. Han (2007) *Cell* **128**, 295-308.
341. Mechanism of coupled folding and binding of an intrinsically disordered protein. K. Sugase, H.J. Dyson and P.E. Wright (2007), *Nature* **447**, 1021-1027. (*News & Views by D. Eliezer and A.G. Palmer in Nature 447, 920, 2007; Faculty of 1000 Exceptional Paper*).
342. Embryonic neural inducing factor Churchill is not a DNA-binding zinc finger protein: Solution structure reveals a solvent-exposed β -sheet and zinc binuclear cluster. B.M. Lee, B.A. Buck-Koehntop, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2007), *J. Mol. Biol.* **371**, 1274-1289. (*Faculty of 1000 recommended paper, January, 2008*).
343. Structure of the Wilms tumor suppressor protein zinc finger domain bound to DNA. R. Stoll, B.M. Lee, E.W. Debler, J.H. Laity, I.A. Wilson, H.J. Dyson and P.E. Wright (2007), *J. Mol. Biol.* **372**, 1227-1245.
344. Tailoring relaxation dispersion experiments for fast-associating protein complexes. K. Sugase, J.C. Lansing, H.J. Dyson and P.E. Wright (2007) *J. Amer. Chem. Soc.* **129**, 13406-13407.
345. Overexpression of post-translationally modified peptides in *Escherichia coli* by co-expression with modifying enzymes. K. Sugase, M.A. Landes, P.E. Wright, M.A. Martinez-Yamout (2008) *Protein Expr. Purif.* **57**, 108-115.
346. Structural characterization of partially folded intermediates of apomyoglobin H64F. S. Schwarzinger, R. Mohana-Borges, G.J.A. Kroon, H.J. Dyson and P.E. Wright (2008) *Protein Sci.* **17**, 313-321.
347. NMR relaxation study of the complex formed between CBP and the activation domain of the nuclear hormone receptor coactivator ACTR. M-O. Ebert, S-H. Bae, H.J. Dyson and P.E. Wright (2008) *Biochemistry* **47**, 1299-1308.
348. Modeling transient collapsed states of an unfolded protein to provide insights into early folding events. D.J. Felitsky, M.A. Lietzow, H.J. Dyson and P.E. Wright (2008) *Proc. Natl Acad. Sci., USA* **105**, 6278-6283.
349. The kinetic and equilibrium molten globule intermediates of apoleghemoglobin differ in structure. C. Nishimura, H.J. Dyson and P.E. Wright (2008) *J. Mol. Biol.* **378**, 715-725.
350. How do proteins interact? D.D. Boehr and P.E. Wright (2008) *Science* **320**:1429-1430.
351. Conformational relaxation following hydride transfer plays a limiting role in dihydrofolate reductase catalysis. D.D. Boehr, H.J. Dyson and P.E. Wright (2008) *Biochemistry* **47**, 9227-9233.
352. Hierarchical folding mechanism of apomyoglobin revealed by ultra-fast H/D exchange coupled with 2D NMR. T. Uzawa, C. Nishimura, S. Akiyama, K. Ishimori, S. Takahashi, H.J. Dyson and P.E. Wright (2008) *Proc. Natl Acad. Sci., USA* **105**, 13859-13864.

353. The intrinsically disordered RNR inhibitor Sml1 is a dynamic dimer. J. Danielsson, L. Liljedahl, E. Bárány-Wallje, P. Sønderby, L.H. Kristensen, M.A. Martinez-Yamout, H.J. Dyson, P.E. Wright, F.M. Poulsen, L. Måler, A. Gråslund and B.B. Kragelund (2008) *Biochemistry* **47**, 13428-13437.
354. Linking folding and binding. P.E. Wright and H.J. Dyson (2009) *Current Opinion in Structural Biology*, **19**, 31-38.
355. Mapping the interactions of the p53 transactivation domain with the KIX domain of CBP. C.W. Lee, M. Arai, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2009) *Biochemistry* **48**, 2115-2124.
356. Functional unfolded proteins: How, when, where, and why? H.J. Dyson, S-C Sue and P.E. Wright (2009). **In:** Water and Biomolecules – Physical Chemistry of Life Phenomena (K. Kuwajima et al., Eds.), Springer-Verlag Berlin Heidelberg, pp. 123-136.
357. Mapping protein folding landscapes by NMR relaxation P.E. Wright, D.J. Felitsky, K. Sugase and H.J. Dyson (2009). **In:** Water and Biomolecules – Physical Chemistry of Life Phenomena (K. Kuwajima, et al., Eds.), Springer-Verlag Berlin Heidelberg, pp. 1-11.
358. Structural basis for recruitment of CBP/p300 coactivators by STAT1 and STAT2 transactivation domains. J.M. Wojciak, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2009) *EMBO J.* **28**, 948-958.
359. Cooperative regulation of p53 by modulation of ternary complex formation with CBP/p300 and HDM2. J.C. Ferreon, C.W. Lee, M. Arai, M.A. Martinez-Yamout, H.J. Dyson and P.E. Wright (2009) *Proc. Natl Acad. Sci., USA* **106**, 6591-6596.
360. Prediction of the Rotational Tumbling Time for Proteins with Disordered Segments S.-H. Bae, H.J. Dyson and P.E. Wright (2009), *J. Amer. Chem. Soc.* **131**, 6814-6821